

Spondylolisthesis: Why Exercise Helps

By PT Ahmad Al Hariri

A Clinical Guide to Conservative Rehabilitation and Movement

What Is Spondylolisthesis?

Spondylolisthesis describes the abnormal anterior (forward) displacement of one spinal vertebra relative to the vertebra immediately below it. This condition most commonly occurs in the lower lumbar spine, particularly at the L4-L5 or L5-S1 levels. It is fundamentally different from a herniated or "slipped" disc, although both conditions can produce similar back or leg symptoms.

From a physical therapy standpoint, the underlying cause—whether it stems from degenerative changes, a defect in the pars interarticularis (isthmic), developmental factors, or mechanical stress—is important. However, our primary clinical focus is on the patient's presentation: their symptoms, movement tolerance, and functional limitations. Rehabilitation is not about altering structural anatomy, but rather optimizing function, improving strength, and managing symptom triggers.




Figure 1: Illustration of anterior translation (spondylolisthesis) of the L4 vertebra over L5.

Disclaimer: This document is intended to guide conservative physical therapy management. It does not replace medical consultation, nor does it advise on surgical interventions or medication. Patients should always consult a physician for diagnostic imaging and medical clearance.

Clinical Presentation and Symptoms

The presentation of spondylolisthesis is highly variable. Some individuals have a significant radiographic slip but experience absolutely no symptoms, participating in high-level sports without limitation. Others may experience significant pain and functional impairment.

Common Local Symptoms

- Localized low-back pain, often described as an ache.
- Pain that increases with prolonged standing or walking.
- Perceived stiffness or "giving way" in the lumbar region.
- Discomfort exacerbated by leaning backwards (spinal extension).

Radiating & Neural Symptoms

- Buttock or posterior thigh discomfort.
- Noticeable hamstring tightness or spasm.
- Numbness, tingling, or pain traveling down one or both legs.
- Reduced tolerance for lifting or loaded activities.

Clinical Insight: Imaging vs. Symptoms

The presence of a vertebral slip on an X-ray or MRI does not automatically explain every symptom. As physical therapists, we treat the patient, not just the scan. A careful clinical correlation is required to ensure that the pain is mechanically related to the spinal segment in question, rather than originating from the hip, SI joint, or surrounding soft tissues.

Clinical Assessment Before Starting Exercise

A diagnosis of spondylolisthesis should never be treated with a generic "low back pain" exercise sheet. Before prescribing a routine, a thorough assessment must establish which activities reproduce the patient's symptoms, whether there is true neural compression (radiculopathy or claudication), and what the patient's baseline capacity is.

Key assessments in the clinic include:

- **Symptom Behavior:** Which positions ease the pain? (Often sitting or forward flexion). Which provoke it? (Often standing, walking, or overhead reaching).
- **Neurological Screen:** Testing for measurable weakness, altered reflexes, or sensory deficits in the lower extremities.
- **Mobility and Control:** Observing how the patient moves from sitting to standing, their gait mechanics, and how they stabilize their trunk during limb movement.

Why Exercise Can Help

It is a common misconception that exercise can "push the vertebra back into place." The goal of physical therapy is entirely different. Exercise aims to improve physical capacity, enhance trunk and hip motor control, restore confidence in movement, and increase the body's tolerance for everyday and recreational activities.

By strengthening the musculature surrounding the spine and pelvis, we can reduce excessive mechanical stress on the symptomatic segment, allowing the area to desensitize.




Figure 2: The supine bridge promotes simultaneous core control and hip extensor strengthening without provoking lumbar extension.

Trunk, Core, and Hip Strengthening

Effective rehabilitation focuses on progressive loading rather than permanent avoidance of movement. There is no universal "spondylolisthesis exercise," but programs commonly target the following areas:

- **Abdominal Bracing and Control:** Starting with basic core activation while maintaining normal diaphragmatic breathing. The goal is endurance and control, not rigid breath-holding.
- **Hip Extensor Strengthening:** Exercises like bridging (shown above) or modified squats help shift the mechanical burden of lifting and movement from the lumbar spine to the powerful gluteal and hamstring muscles.
- **Supine and Quadruped Movements:** Marching, heel slides, or bird-dog variations performed in symptom-free ranges to challenge rotary and flexor control.
- **Lower Extremity Strength:** Step-ups, supported squats, and side-lying hip abduction build functional capacity for walking and stair climbing.

Exercise Progression and Modifications

Rehabilitation should follow a staged approach. Increasing the difficulty of an exercise is useful only when the previous level is performed with good technical control and without symptom flare-ups.

1. **Phase 1:** Establish a tolerable starting position and learn basic trunk/pelvic control.
2. **Phase 2:** Increase repetitions or duration to build local muscular endurance.
3. **Phase 3:** Introduce external resistance (bands, weights) and transition to upright, functional movements.
4. **Phase 4:** Reintroduce sport-specific or occupational demands (e.g., dynamic lifting, light running).

What to Modify or Avoid

While we want to encourage movement, certain biomechanical stresses are notorious for irritating spondylolisthesis in the early phases of rehab. Modifications should be individualized, but generally include:

- **End-Range Extension:** Repeated or sustained leaning backwards (such as certain gymnastics poses, heavy overhead pressing, or prone back-bending) often compresses the posterior structures and stretches the anterior restraints. Modify these to neutral-spine activities.
- **High-Impact Loading:** During highly irritable phases, jarring activities (like prolonged running on hard surfaces or jump-landing) may need to be temporarily replaced with cycling, swimming, or brisk walking.
- **Heavy Uncontrolled Flexion:** While forward bending often feels good, heavy lifting with a rounded spine can overload the discs and pars defect. Hinge mechanics (using the hips) must be taught.

When to Consult Your Physician

Physical therapy is highly effective for conservative management, but medical review is critical if you experience: sudden weakness in the legs, loss of bowel or bladder control, severe unrelenting pain that wakes you up at night, or progressively worsening numbness. These require immediate medical evaluation.

Conclusion

A diagnosis of spondylolisthesis does not mean a lifetime of fragility. With a well-structured, progressive physical therapy program focused on core endurance, hip strength, and movement mechanics, the vast majority of patients can successfully manage their symptoms and return to the activities they love. The key is to start at a tolerable level, focus on movement quality, and progressively build capacity over time.